



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

(December 4, 1916) Siskin's (*Spinus pinus*), Acadian Chickadees (*Parus hudsonicus littoralis*), Redpolls (*Acanthis linaria linaria*) American and white-winged Crossbills (*Loxia curvirostra minor* and *L. leucoptera*), Pine Grosbeaks (*Pinicola enucleator leucura*), Evening Grosbeaks (*Hesperiphona vespertina vespertina*), Snow Buntings (*Plectrophenax nivalis nivalis*), Snowy Owl (*Nyctea nyctea*), and Rough-legged Hawks (*Archibuteo lagopus sancti-johannis*) are here, many of them in much greater abundance than for many years.—MRS. GEORGE H. MELLE, *Newton Highlands, Mass.*

**Evening Grosbeak** (*Coccothraustes vespertina vespertina*) **at Cinnaminson, N. J.**—On December 3, 1916, Mr. Charles Evans of Cinnaminson, near Riverton, N. J., found two Evening Grosbeaks feeding on his lawn. He recognized them at once by a colored plate of the species which he had, but supplemented this identification with a direct comparison with the description in Chapman's 'Handbook.' The birds were exceedingly tame and would scarcely get out of his way. Cinnaminson is only eight miles above Philadelphia on the opposite side of the Delaware River and this constitutes the first record of the species south of Plainfield, Fair Haven and Princeton from which places it is recorded in 'Bird-Lore,' for 1911 and 1913. The Cinnaminson birds are therefore the most southern recorded occurrence for this interesting species.—GEORGE SPENCER MORRIS, *Olney, Philadelphia, Pa.*

**Clarence Henry Morrell — A Correction.**—In the 'Ten Year Index to the Auk,' p. XIX, 1915, the birth place of Clarence Henry Morrell is given as Pittsfield, Me., and the date of birth as Feb. 23, 1872. These data were obtained from an obituary notice by Mr. J. M. Swain in 'The Auk' for 1902, p. 423. My attention was recently called to a more extended biographical sketch of Mr. Morrell in the 'Journal of the Maine Ornithological Society,' V, pp. 7-12, 1903, also prepared by Mr. Swain, stating that Mr. Morrell was born at River Hobart, Nova Scotia, Feb. 27, 1872. This statement Mr. Swain advises me is correct and the place and date originally published in 'The Auk' are erroneous.—T. S. PALMER, *Washington, D. C.*